

Instructions For Use FAB-2-IFU

Revision: 2

Rev. Date: Aug. 17, 2014

Page 1 of 2

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

Acid Fast Bacteria (AFB) Stain Kit


Description:

The Acid Fast Bacteria (AFB) Stain Kit is intended for use in the histological visualization of Acid Fast Bacteria and Tubercle Bacilli. This kit is a rapid 15-minute procedure. The lipoid capsule of the acid-fast organism takes up carbol fuchsin and resists decolorization.

Acid Fast Organisms: Bright Red Background: Light Green

- Uses/Limitations: Not to be taken internally. For In-Vitro Diagnostic Use. Histological applications. Do not use if reagents become cloudy. Do not use past expiration date. Use caution when handling reagents. Non-Sterile.
- Control Tissue: Any well fixed paraffin embedded or frozen tissue section cut at 3-5µ. Cell Smear.

Ordering information regarding individual components on back page!



Item #	Kit Contents	Volume	Storage
CFZ030	Carbol Fuchsin Solution	30 ml	18-25°C
AAL030	Acid Alcohol Solution (0.5%)	2 x 30 ml	18-25 <i>°</i> C
LGA030	Light Green Solution	30 ml	18-25 <i>°</i> C

- Precautions: Avoid contact with skin and eyes. Harmful if swallowed. Follow all Federal, State, and local regulations regarding disposal. Use in chemical fume hood whenever possible. Wear protective clothing.
- **Note:** Carbol Fuchsin Solution should be filtered when a thick sheen develops on top of solution. To avoid possible contamination, gloves should be worn when performing this procedure. Do not use tap water prior to application of Carbol Fuchsin Solution as it is reported that Acid Fast Bacteria can be found in some systems. Therefore use of distilled water is recommended whenever possible.

Storage: 18° C



ScyTek Laboratories, Inc. 205 South 600 West Logan, UT 84321 U.S.A.



EC_REP EmergoEurope (31)(0) 70 345-8570 Molsnstraat 15 2513 BH Hague, The Netherlands

Doc: IFU-Template18-25rev2



Instructions For Use FAB-2-IFU

Revision: 2

Rev. Date: Aug. 17, 2014

Page 2 of 2

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Tel. (435) 755-9848 - Fax (435) 755-0015 - www.scytek.com

Procedure (Standard):


- 1. Deparaffinize sections if necessary and hydrate to distilled water.
- 2. Incubate slide in Carbol Fuchsin Solution for 15 minutes.
- 3. Rinse for 2 minutes in running tap water followed by 2 changes of distilled water.
- 4. Decolorize in Acid Alcohol Solution (0.5%) until sections are a pale pink color.
- 5. Dip slide twice in distilled water to rinse.
- 6. Counterstain in Light Green Solution for 1-2 minutes.
- 7. Rinse in several changes of distilled water.
- 8. Dehydrate quickly through graded alcohols ending with 2 changes in absolute alcohol.
- 9. Clear, and mount in synthetic resin.

References:

- 1. Sheenan, D.C., Hrapchak, B.B. Theory and Practice of Histotechnology, 2nd Edition. Battelle Press, Columbus, OH.
- 2. Carson, F.L., 1996, Histotechnology; A Self-Instructional Text, 2nd Edition. ASCP Press, Chicago, IL.

Bulk Reagent Ordering Information and Current Pricing at www.scytek.com

Description:	Catalog #	Volume
Carbol Fuchsin Solution	CFZ125 CFZ500 CFZ999	125 ml 500 ml 1000 ml
Acid Alcohol Solution (0.5%)	AAL500 AAL999	500 ml 1000 ml
Light Green Solution	LGA125 LGA500 LGA999 LGA3800	125 ml 500 ml 1000 ml 1 Gallon









EC REP EmergoEurope (31)(0) 70 345-8570 Molsnstraat 15 2513 BH Hague, The Netherlands

Doc: IFU-Template18-25rev2